

*ikṣvāku* = *kaṭutumbī* = *tiktatumbī*

- (1) a) LAGENARIA SICERARIA STANLEY = L. VULGARIS SERINGE (Avk; Bpn, p. 671; Dgv, nr. 165; DWH 2, p. 63 : the bitter variety; Gul; KB 2, p. 1116-1119; Nadk. 1, nr. 1419; V 2, p. 28) = CUCURBITA LAGENARIA LINN. (Avk; Kav; KB 2, p. 1116-1119; Nadk. 1, nr. 1419; V 2, p. 28) = C. SICERARIA MOLINA;
- b) the wild variety of a) (MW; PW; Roxb., p. 700; Vss);
- (2) CUCURBITA PEPO LINN. (Nadk. 1, nr. 1419);
- (3) CITRULLUS COLOCYNTHIS SCHRAD. (MW; PW).

*indrayava*

= the seeds of *kuṭaja*.

*utpala*

- (1) NYMPHAEA STELLATA WILLD. (Chopra; Dutt; Gul; HK; KB 1, p. 113-115; Na 1, nr. 18; Nadk. 1, nr. 1726) = N. CYANEA ROXB. (HK; PW) = N. CAERULEA SAVIGN. (Kav; MW) [= N. STELLATA LINN.] (Kav) = CASTALIA STELLATA BLUME (KB 1, p. 113-115);
- (2) N. LOTUS LINN. (Nadk. 1, nr. 1726) [= N. LOTUS HOOK.F. ET THOMS.] (KB 1, p. 112-113) = N. RUBRA ROXB. EX SALISB. (KB 1, p. 112-113) = N. RUBRA (Nadk. 1, nr. 1726; PW);
- (3) N. ODORATA [SOLAND. IN] AIT. (Ainslie 2, p. 381);
- (4) COSTUS SPECIOSUS Sm. (MW; PW, = *kuṣṭha*; Vss : = *kuṣṭha*);
- (5) SAUSSUREA LAPPA C. B. CLARKE (Nadk. 1, nr. 2239, = *kuṣṭha*);
- (6) = *padmakāṣṭha* (Vss);
- (7) BOMBAX CEIBA LINN. = B. MALABARICUM DC. = B. HEPTAPHYLLUM Cav. (Vss).

*udīcyā* = *udīcī*

- (1) = *bālā* (HK; Kav; KB 1, p. 323-324);
- (2) = *hrīvera* (KB 1, p. 323-324 : = *bālā* = *hrīvera*; Vss);
- (3) a kind of perfume (MW; PW); the term *udīcyā*, though occurring thirteen times in Ca., is not explained by Cakra.

*udumbara*

- (1) FICUS RACEMOSA WALL. = F. RACEMOSA (Nadk. 1, nr. 1041) [= F. RACEMOSA LINN.] (acc. to Chopra, but the Index Kewensis indicates