

- c) *vaijayantītulasī* (Vss, unidentified);
- d) = *barbarika* = *ajagandhikā* : a kind of Ocimum (PW);
- (4) CLERODENDRUM SERRATUM SPRENG. (KB 3, p. 1948-1949);
- (5) C. SIPHONANTHUS [R. BR. IN] AIT. (MW; PW);
- (6) GYNANDROPSIS PENTAPHYLLA DC. = CLEOME PENTAPHYLLA LINN. (MW; PW);
- (7) CLEOME VIScosa LINN. (KB 1, p. 183-185);
- (8) ACACIA ARABICA WILLD. (Nadk. 1, nr. 12) = MIMOSA ARABICA LAM. (Roxb., p. 421);
- (9) A. FERRUGINEA DC. (Nadk. 1, nr. 12);
- (10) A. LEUCOPHLOEA WILLD. (KB 2, p. 924-925 : *svetabarbara*);
- (11) = *pītacandana* (see *candana*) (PW; Vss);
- (12) a partic. fragrant plant (MW);
- (13) [unguis odoratus] (MW; PW : = *vyāghranakha* : a) possibly *οὐνξ* of Dioscorides, unguis odoratus; see Dietz, p. 57; b) EUPHORBIA ANTIQUORUM LINN. = TITHYMALUS ANTIQUORUM MOENCH.).

barbaracandana

Dn, *varga* 3, 9 : *barbarika* is a white, odourless type of *candana*; Rn, *varga*, 12, 17: *barbaraka*, (also called) white *barbaraka*, is (a kind of *candana* that is) fragrant; a synonym (see Dn and Rn) is *barbarottha*, explained in PW as white sandalwood.

barbarikā

Is not recorded in the dictionaries; see *vanabarbarikā*.

barbarī

- (1) a) OCIMUM BASILICUM LINN. (Avk; Bpn, p. 511-512; KB 3, p. 1961-1964)[= O. BASILICUM VAR. THYRSIFLORA BENTH.] (Avk); see Hooker 4, p. 608;
- b) a kind of Ocimum, = *barbara* (MW; PW);
- c) Bpn, p. 511 : = *parṇāsa*; three types are distinguished : a) black : *kūtheraka*, b) white : *arjaka*, c) with leaves like *vatā* : *vatāpatra*;
- (2) ORTHOSIPHON GRANDIFLORUS BOLDINGH EX HEYNE¹ (Bpn, p. 511-512).