

kr̥mīśatru

Probably = vidāṅga: Embelia ribes Burm.f. (Kai.Nigh., ośadhivarga 1147-1148: vidāṅga = kr̥mījit, kr̥mihantr, kr̥mihara, kr̥mihrt, jantughna); however, kr̥mighna is also a synonym of kolakanda, palāṅdu, pāribhadra, bījapūrṇa and bhallātaka.

ketakī

Pandanus tectorius Soland. (Dgv, nr.333; KB 4, p.2592-2593: ketaka) = P. odoratissimus Linn.f. (WIRM 7, p.218) (P. odoratissimus Roxb. acc. to Kai.Nigh., ośadhivarga 1483-1485 and Bpn, puṣpavarga 42, P. odora-tissimus Willd. acc. to Nadk. 1, nr.1785, Raison and DEP 6, p.5).

kovidāra

(1) Bauhinia variegata Linn. (HK; Dgv, nr.121; Gul; Bpn, guḍūcyādivarga 101-102; Nadk. 1, nr.331; Na 1, nr.180; Chopra; Bt.; KB 2, p.898-900);

(2) B. purpurea Linn. (Bpn, guḍūcyādivarga 101-102; Bt.; V. 3, p.24; Kai.Nigh., ośadhivarga 933; Sharma; KB 2, p.897-898);

(3) B. tomentosa Linn. (Bpn, guḍūcyādivarga 101-102).

kośātaka, kośātakī

(1) a) Luffa acutangula Roxb. (HK; Gul; Nadk. 1, nr.1514; Na 1, nr.230; WIRM 6, p.177; Sharma; Bt.; Chopra; Dutt; KB 2, p.1121-1123; Kai.Nigh., ośadhivarga 568-569);

b) L. acutangula Roxb. var. amara (Roxb.) C.B. Clarke (DWH 2, p.80-81; KB 2, p.1123-1125);

(2) L. echinata Roxb. (Nadk. 1, nr.1517; Bt.);

(3) L. graveolens Roxb. (Nadk. 1, nr.1518);

(4) L. cylindrica (Linn.) M.J. Roemer (Bt.).

kośāmra

Schleichera oleosa (Lour.) Oken = S. trijuga Willd. (Bpn, āmrādivarga 23-24; Nadk. 1, nr.118; KB 1, p.629-631; V. 3, p.29) [= S.