

Hooker as a synonym; described in Roxb., p.555; P. sublobatus Roxb. = P. trinervius Heyne acc. to Cooke 1,p.402 and Gamble 1,p.256);

(3) Vigna dalzelliana (Kuntze) Verdcourt = Phaseolus dalzelli Cooke (PA 1,7,p.72-73) = P. pauciflorus Dalz. non G. Don nec Benth. (PA 1,7, p.72-73: P. pauciflorus Dalz.; see WIRM 10,p.475 on the nomenclature);

(4) Vigna umbellata (Thunb.) Ohwi et Ohashi = Dolichos umbellatus Thunb. = Phaseolus calcaratus Roxb. (PA 1,7,p.66; see WIRM 10,p.496 on the nomenclature);

(5) Vigna trilobata (Linn.) Verdcourt = Dolichos trilobatus Linn. = Phaseolus trilobus sensu Ait. et auct. (DGV 5,p.227-228: P. trilobus Ait.; see WIRM 10,p.495 on the nomenclature);

(6) Vigna mungo (Linn.) Hepper = Phaseolus mungo Linn. (DGV 2-3,p. 346; P. mungo Linn. is the correct name acc. to Lily M. Perry, but see WIRM 10,p.476 on the nomenclature);

(7) Atylosia goensis Dalz. (Chopra; PA 1,7,p.77-78) = A. barbata Baker (Nadk. 1,nr.286; PA 1,7,p.77-78) = Dunbaria barbata Benth. (PA 1,7,p.77-78);

(8) Pueraria phaseoloides Benth. (PA 1,7,p.66);

(9) Calopogonium mucunoides Desv. (PA 1,7,p.66).

vandā

(1) Dendrophthoe falcata (Linn.f.) Ettingshausen (Bpn, guḍūcyādivarga 261; Chopra; WIRM 3,p.34) = Loranthus falcatus Linn.f. (Chopra; KB 3,p.2180-2181; WIRM 3,p.34) = L. longiflorus Desr. (Bpn, guḍūcyādivarga 261; Chopra; KB 3,p.2180-2181; DEP 5,p.92-93; WIRM 3,p.34);

(2) Vanda tessellata Lodd. ex Loud. = V. roxburghii R.Br. = Epidendrum tessellatum Roxb. (MW; Vśś);

(3) Viscum album Linn. (Bpn, guḍūcyādivarga 261);

(4) V. articulatum Burm.f. (KB 3,p.2184-2185).

hamsapādī

(1) Cayratia pedata (Wall.) Gagnep. = Vitis pedata Vahl ex Wall. (HK) = Cissus pedata Lam. (HK);

(2) Adiantum philippense Linn. = A. lunulatum Burm. (Bpn, guḍūcyādivarga 256; Bt.; Dgv, nr.348; Gul.; Kai.Nigh., oṣādhivarga 766-767; KB 4,p.2735-2736; Na 2,nr.567);