

- (4) MANGIFERA INDICA LINN. (Vśś);  
 (5) SPONDIAS MANGIFERA WILLD. (Vśś, = *mācīkā* = *āmrātaka*);  
 (6) PHYLLANTHUS NIRURI LINN. (KB 3, p. 2225-2227);  
 (7) = *palāśīlatā* (PW; Vśś); acc. to KB 1, p. 788-789 and Nadk. 1, nr. 397 *palāśīlatā* is BUTEA SUPERBA ROXB.;  
 (8) Kav (footnote p. 352) : "Cakra says (on Ca.Sū. 27, 121) that it is a little shrub growing in Kāmarūpa and its neighbourhood; Gaṅgādhara believes that *amlīkā* is otherwise called *amlāṛdraka*, a sour variety of ginger; the plant is difficult to identify."

*amlīkā*

- (1) TAMARINDUS INDICA LINN. (HK);  
 (2) OXALIS CORNICULATA LINN. (MW);  
 (3) = *amlīkā* (PW).

*arjaka*

- (1) OCIMUM GRATISSIMUM LINN. (MW; PW; V 1, p. 78);  
 (2) O. BASILICUM LINN. (Avk; Bpn, p. 511-512);  
 (3) O. SANCTUM LINN. (KB 3, p. 1965-1968) = O. VILLOSUM ROXB. (Roxb., p. 462);  
 (4) O. CANUM SIMS (Chopra; KB 3, p. 1960-1961, = *kṣudratulasī*) = O. AMERICANUM LINN. (Chopra);  
 (5) ORTHOSIPHON GRANDIFLORUS BOLDINGH EX HEYNE (Bpn, p. 511-512).

*aśvattha*

FICUS RELIGIOSA LINN. = UROSTIGMA RELIGIOSUM GASP.

*āḍhakī*

CAJANUS CAJAN (LINN.) MILLSP. [Intern. Code] (Chopra; IRM 2, p. 6-11) = C. INDICUS SPRENG. (Avk; Bpn, p. 647; Chopra; Dutt; DWH 1, p. 489; Gul; HK; IRM 2, p. 6; Kav; KB 1, p. 809-811; MW; Na 1, nr. 135; Nadk. 1, nr. 409; PW; V 1, p. 82; Vśś; Watt CP, p. 196) = C. BICOLOR DC. (Avk) = C. FLAVUS DC. (Avk) = CAJAN CAJAN MILLSP. = CYTISUS CAJAN LINN. (Avk; Dutt; HK; Kav; V 1, p. 82; Vśś); there are two varieties of this plant, var. flavus DC. (corolla yellow) and var. bicolor DC. (corolla veined with red), both cultivated in India; these varieties have been looked upon as species by some authors; see Watt CP, p. 196-197 and Purseglove 1, p. 237.