

kañkola

- (1) PIPER CUBEBA LINN.F. (Bpn, p. 258-259; Dgv, nr. 267; DWH 3, p. 180-183; HK; MW; PW; V 2, p. 66);
 (2) a particular substance sold under that name (Vśś).

kañgu

- (1) SETARIA ITALICA BEAUV. (Avk; Chopra; Dutt; DWH 3, p. 619; HK; KB 4, p. 2718-2719; Nadk. 1, nr. 2291; V 2, p. 113; Watt CP, p. 988-989) = PANICUM ITALICUM LINN. (Ainslie 1, p. 226-227; Avk; Chopra; Dutt; Gul; HK; Nadk. 1, nr. 1793; PW; Roxb., p. 101-102; Vśś; Watt CP, p. 988-989);
 (2) PANICUM FRUMENTACEUM (Nadk. 1, nr. 1791); Nadk. considers P. FRUM. as a synonym of SETARIA ITALICA BEAUV., but this is not confirmed by the Index Kewensis, which distinguishes two species : P. COLONUM LINN. = P. FRUMENTACEUM ROXB., and SORHGUM VULGARE PERS. = P. FRUMENTACEUM SALISB.; Hooker (7, p. 30-32) regards P. FRUMENTACEUM ROXB. as a synonym of P. CRUS-GALLI LINN. VAR. FRUMENTACEUM and describes P. COLONUM LINN. as a separate species; Cooke (3, p. 446-447) says that P. FRUMENTACEUM ROXB. is a synonym of P. STAGNUM RETZ. VAR. FRUMENTACEA TRIM.;
 (3) PAPAVER DUBIUM (Bpn, p. 653); the Index Kewensis mentions two plants of this name : P. DUBIUM LINN., and P. DUBIUM TENORE = P. PINNATIFIDUM MORIS.; it may be that P. D. LINN. is meant, because this species is also described in the Flora Iranica.

kaṭukā

- (1) PICRORRHIZA KURROA ROYLE EX BENTH. (Bpn, p. 70; Chopra) = P. KURROA ROYLE (DWH 2, p. 459; Dy, p. 577-578) = [P. KURROA BENTH.] (Dutt; HK; KB 3, p. 1825-1826; Nadk. 1, nr. 1907) = P. KURROA (Avk; Dgv, nr. 235; HB; Sanyal; Vśś);
 (2) = *kaṭvī* (PW : name of several plants);
 (3) = *kaṭurohiṇī* (PW);
 (4) = *tāmbūlī* (PW; acc. to PW : ARECA CATECHU LINN., but acc. to PA 1, 9, p. 71 : PIPER BETLE LINN.);
 (5) RUELLIA LONGIFOLIA (PW: = *kulīkavṛkṣa*); the Index Kewensis enumerates five species of this name; the genus Ruellia consists of herbs and undershrubs (Hooker 4, p. 411), not of trees; the species meant by PW may be R. LONGIFOLIA T. ANDERS., described by Hooker (4, p. 412);