

(3) SACCHARUM SPONTANEUM LINN. (Vśś : = *kāśā*). Many authors equate *darbha* with *kuśā*, but Ḍa. (on Su.Sū. 38, 75) distinguishes them clearly : “*kuśā* is short and soft and has leaves with the form of a needle; *darbha* has leaves which are broad, long and rough.”

*dāḍīma*

PUNICA GRANATUM LINN.

*dāruharidrā*

(1) BERBERIS ARISTATA DC. (AFM 1, p. 60; Avk; Bpn, p. 119; Chopra; Dgm, p. 44; Dgv, nr. 17; Dutt; DWH 1, p. 64-68; Dy, p. 35-37; Ghosh; Gul; HK; IRM 1, p. 178; KB 1, p. 102-103; Nadk. 1, nr. 335; PA 1, 2, p. 15; V 5, p. 65) [= B. ARISTATA VAR. FLORIBUNDA HOOK.F. ET THOMS.] (KB 1, p. 102-103) = B. CORIARIA ROYLE EX LINDL. (KB 1, p. 102-103) = B. CHITRIA D. DON (Dgm, p. 44);

(2) B. ASIATICA ROXB. EX DC. (Avk; Bpn, p. 119; Dgm, p. 44-45; Dutt; DWH 1, p. 64-68; Dy, p. 35-37; HB; Kav; KB 1, p. 105-106; Na 1, nr. 16; PA 1, 2, p. 15; Sanyal; Vśś);

(3) B. LYCIUM ROYLE (Bpn, p. 119; Dgm, p. 44-45; DWH 1, p. 64-68; Dy, p. 35-37);

(4) B. VULGARIS LINN. (Avk; Dgm, p. 44-45; PA 1, 2, p. 15);

(5) B. CORIACEA BRANDIS EX GAMBLE (Dgm, p. 44-45);

(6) B. NEPALENSIS (Dgm, p. 44-45); the Index Kewensis indicates two species : a) B. NEPALENSIS SPRENG.; b) B. VULGARIS LINN. = B. NEPALENSIS HORT. EX C. KOCH;

(7) [COSCIINIUM FENESTRATUM (GAERTN.) COLEBR.] (Chopra; IRM 2, p. 360; KB 1, p. 84-86) = C. FENESTRATUM COLEBR. (AFM 1, p. 60; Bpn, p. 121-122 : used as a substitute; Nadk. 1, nr. 685; PA 1, 2, p. 15; V 5, p. 71; Vśś) = MENISPERMUM FENESTRATUM GAERTN. (KB 1, p. 84-86; PA 1, 2, p. 15);

(8) CURCUMA AROMATICA SALISB. (MW) (usually called *vanaharidrā*);

(9) C. XANTHORRHIZA ROXB. (MW);

(10) MORINDA UMBELLATA LINN. (KB 2, p. 1296-1297).

*dārvī*

= *dāruharidrā*.