

V 6, p. 89) = GARDENIA ULIGINOSA RETZ. (KB 2, p. 1272-1273) = POSOQUERIA ULIGINOSA (WILLD.) ROXB. (KB 2, p. 1272-1273);

(4) TINOSPORA CORDIFOLIA MIERS = COCCULUS CORDIFOLIUS DC. (PW : = *kandaguḍūcī*; Vśś : = *kandaguḍūcī*; identical with *guḍūcī*).

*pippalī*

PIPER LONGUM LINN.

*pīlu*

(1) SALVADORA PERSICA LINN. (Avk; Chopra : *brhatpīlu*; Dgv, nr. 209; Dutt; DWH 2, p. 380-384; Gul; HB; HK; Kav; KB 2, p. 1537-1539; MW; Nadk. 1, nr. 2203; PW; Vśś) = S. INDICA WIGHT (Chopra; Dutt; HK; Kav; KB 2, p. 1537-1539; Vśś) = S. WIGHTIANA PLANCH. EX THW. (Nadk. 1, nr. 2203) = S. OLEOIDES DECNE. (Bpn, p. 577; Chopra; DWH 2, p. 380-384; KB 2, p. 1539-1540; Vśś); acc. to the Index Kewensis S. PERSICA LINN. = S. OLEOIDES DECNE.; Chopra considers them to be two separate species; so do Kirtikar and Basu; Hooker says in his Flora of India 3, p. 620, that S. OLEOIDES DECNE. is closely allied to S. PERSICA LINN., but well distinguished by Brandis (see Brandis' Indian Trees, p. 453): the Flora Iranica also regards S. PERSICA and S. OLEOIDES as separate species;

(2) CAREYA ARBOREA ROXB. (KB 2, p. 1061-1063; MW; PW).

*punarnavā*

(1) BOERHAAVIA REPENS LINN. (Chopra; Dgm, p. 210; DWH 3, p. 130-132; Gul; HK; IRM 1, p. 199; Nadk. 1, nr. 363; PA 1, 5, p. 9-26; Sanyal) = B. DIFFUSA LINN. (Ainslie 2, p. 205-207; Avk; Bpn, p. 421 : *śvetapunarnavā*; Chopra; Dgv, nr. 258; Dutt; Dy, p. 655-656; HB; HK; IRM 1, p. 199; Kav; KB 3, p. 2045-2048; Nadk 1, nr. 363; PA 1, 5, p. 9-26; V 6, p. 125; Vśś; Wise) [= B. PROCUMBENS ROXB.] (KB 3, p. 2045-2048; Nadk. 1, nr. 363; PA 1, 5, p. 9-26; Roxb., p. 49) [= B. ERECTA BANKS EX ROXB.] (HK); unclear are :

- a) B. PROCUMBENS, mentioned by Dutt, Nadk. (1, nr. 363) and Vśś, which may be the plant designated as B. PROCUMBENS ROXB. by some authors, or another species: B. PROCUMBENS BANKS EX ROXB.;
- b) B. ERECTA, mentioned by Dutt, Nadk. (1, nr. 363) and Vśś, which may be B. ERECTA ELL., a synonym of B. REPENS LINN., or a different species : B. ERECTA LINN. = B. DIANDRA BURM.F.;